

AD-A140 114

GENETIC AND PHYSICAL STRUCTURES OF HYBRID BACTERIOPHAGE 1/1  
GENOMES(U) HAHNEMANN MEDICAL COLL AND HOSPITAL  
PHILADELPHIA PA N YAMAMOTO JAN 82 DAMD17-79-C-9134

UNCLASSIFIED

F/G 6/13

NL


END

DATE

FORMED

5 84

OTIC



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

①

AD \_\_\_\_\_

ANNUAL PROGRESS REPORT

Genetic and Physical Structures of  
Hybrid Bacteriophage Genomes

Annual Progress Report

Nobuto Yamamoto, Ph.D.

January, 1982

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND,  
Fort Detrick, Frederick, Maryland 21701

Contract No. DAMD 17-79-C-9134

Hahnemann Medical College  
Philadelphia, Pennsylvania 19102

Approved for public release;  
Distribution unlimited

The findings in this report are not to be construed as an official  
Department of the Army position unless so designated by other  
authorized documents.

DTIC  
ELECTE  
APR 15 1984  
S E D

AD A140114

DTIC FILE COPY

84 04 18 128

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A14011	4
4. TITLE (and Subtitle) Genetic and Physical Structures of Hybrid Bacteriophage Genomes	5. TYPE OF REPORT & PERIOD COVERED Annual Progress Report (From 2/1/81 to 1/31/82)	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s) Nobuto Yamamoto	8. CONTRACT OR GRANT NUMBER(s) DAMD 17-79-C-9134	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Hahnemann Medical College 230 North Broad Street Philadelphia, PA 19102	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102A.3M161102BS10.AE.062	
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Medical Research and Development Command, Fort Detrick, Frederick, MD 21701	12. REPORT DATE January 1982	
	13. NUMBER OF PAGES 13	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bacteriophage, Hybrid bacteriophage, <u>Salmonella typhimurium</u> , Hybrid bacteria, <u>E. coli-S. typhimurium</u> hybrid, Genetic homology, Genetic recombination, Antigen conversion, Bacteriophage tail, Mutation, and Transduction. <u>ESCHERICHIA</u> <u>SALMONELLA</u> <u>lambda</u>		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <u>E. coli-S. typhimurium</u> hybrids provides excellent systems to isolate bacteriophage hybrids between <u>Salmonella</u> phage P22 and <u>E. coli</u> phages such as $\lambda$ , $\phi$ 80 and Mu-1. A variety of hybrid types such as $\lambda$ -P22Im $\lambda$ C, $\phi$ 80ImmP22, and MuImmP22 have been isolated and characterized. Studies of these hybrid phages provide invaluable information for the genetic evolution, mutation, transduction, immunity controls and gene expression of bacteriophages.		

*phi*  
 We have isolated  $\phi 80immP22dis$  and  $MuimmP22dis$  hybrid phages carrying both c and Im regions of P22. Derivatives of  $\phi 80immP22dis$  which have lost the dis function were recovered from  $\phi 80immP22dis$  lysogens. Such dis derivatives were formed by replacement of a P22 phage segment containing att through Im regions with bacterial regions adjacent to prophage insertion site. Therefore the dis hybrid derivatives are high frequency transducing phages for proA, argF and metD but not for tryptophan.

*phi*  
 Although homology between  $\phi 80immP22$  hybrids and P22 have been mapped in detail,  $\phi 80$  segments in the hybrids have not been analyzed. Mapping of  $\phi 80$  segments are now feasible since we found WR4027 amber suppressors for  $\phi 80$  amber mutants.

P22- $\lambda$  hybrid phage carries a segment of the  $\lambda$  early regions and the entire late region of P22. Thus the P22 late genes should be regulated by the  $\lambda$  early genes. This was analyzed by complementation capacity of  $\lambda$  or  $\phi 80$  for P22 amber mutants within P22 early regions. P22 amber mutants: both am24 for early regulatory gene and am23 for late regulatory gene and am7 for endolysin gene were complemented by  $\lambda$  and  $\phi 80$  phages. However am12 for control of P22 DNA replication gene was not complemented.

Salmonella phages P22 and ES18 are serologically and morphologically unrelated. However ES18 has a homology with the entire P22 early regions and the gene for generalized transduction, suggesting that ES18 is a generalized transducing hybrid phage as a consequence of recombination between P22 like group A Salmonella phage and a group B phage.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



## CONTENTS

	Page
Front cover	1
Report document page with abstract (DD Form 1473)	2
Cover for progress report	4
Summary	5
Foreword	7
Progress	8
1. Isolation of a new hybrid between Salmonella phage P22 and coli mutator phage 8	8
2. Isolation of amber suppressor strains of <u>E. coli-S. typhimurium</u> hybrid WR4028/22 met <sup>+</sup>	8
3. Isolation and characterization of transducing $\phi$ 80immP22 hybrids	9
4. Complementation of the P22 early gene functions with $\lambda$ and with $\phi$ 80.	10
5. A new isolation procedure for $\lambda$ -P22Im $\lambda$ C hybrids.	10
6. Homology between two generalized Salmonella transducing phages ES18 and P22.	11
Publications	12
Distribution list	13

## SUMMARY

Replication of P22 phage in Mu-1 lysogen gave rise to a hybrid phage designated MuimmP22 carrying the protein coat of Mu-1 and c region of P22. Since P22 carries antirepressor gene, P22 can grow in MuimmP22 lysogen and gives rise to another hybrid type designated MuimmP22dis which carries both c and Im region of P22.

We have also isolated  $\phi$ 80immP22dis hybrids which carries both the immunity regions (c and Im) and prophage attachment region (att) of P22. Hybrid derivatives which have lost the dis function (i.e.  $\phi$ 80immP22dis<sup>-</sup>) have been recovered from  $\phi$ 80immP22dis lysogens. These dis<sup>-</sup> derivatives were formed by replacement of a P22 phage segment containing att through Im regions with bacterial segment adjacent to the prophage insertion site. Thus these dis<sup>-</sup> hybrid derivatives are high frequency transducing phages for proline but not tryptophan. They cotransduce proA, argF and metD genes.

We have mapped homologous segment between  $\phi$ 80immP22 hybrids and P22 in detail. To analyse the  $\phi$ 80 segments in the hybrids, we isolated WR4027 amber suppressors for  $\phi$ 80 amber mutants.

P22- $\lambda$  hybrid phage carries a segment of the  $\lambda$  early regions and the entire late region of P22. Thus the  $\lambda$  early genes should regulate expression of the P22 late genes in P22- $\lambda$  hybrid phage. This was analyzed by complementation capacity of  $\lambda$  or  $\phi$ 80 for P22 amber mutants within their early regions. Complementation should be demonstrable by superinfecting  $\lambda$  or  $\phi$ 80 lysogens with P22 amber mutants because the P22 ant function induces these prophages. P22 amber mutants; both am24 for early regulatory gene and am23 for late regulatory genes, and am7

for endolysin gene were complemented by  $\lambda$  and  $\phi$ 80 phages. However, am12 for control of DNA replication gene was not complemented.

Although two generalized transducing Salmonella phages P22 and ES18 are serologically and morphologically unrelated. ES18 has a homology with the entire early regions and gene for generalized transduction of P22, suggesting that the generalized transducing phage ES18 is created as a consequence of recombination between a group A phage (P22 like phage) and a group B phage.



## FOREWORD

Fundamental studies of bacterial and viral genetics not only play an important role in increasing our knowledge of the action of viruses in disease processes, but also contribute greatly to our knowledge of the whole problem of cell replication, genetic transfer, gene control, morphogenesis, and antigen conversion. The significance of the study of bacterial hybrids between E. coli and Salmonella has greatly broadened with the discoveries of hybrid phages between coliphage and Salmonella phage. The study supported by this contract will bring many important answers for mechanisms of genetic evolution, transduction, recombination, gene expression, antigen conversion, morphogenesis and viral replication. In addition, such newly constructed hybrids may prove useful in achieving intergeneric transduction via a hybrid phage vector of chromosomal genes from different genera of enterobacteriace. Therefore, such hybrid phages may serve as useful vectors in the genetic engineering of a polyvalent oral attenuated vaccine which expresses immunogenic determinants for antigens of Shigella, Salmonella, and perhaps even cholera.

## PROGRESS

### 1. Isolation of a new hybrid between Salmonella phage P22 and coli mutator phage Mu-1

Phage Mu-1 is unable to grow in a smooth E. coli-S. typhimurium hybrid strain WR4028 but able to grow in a rough strain WR4027. In contrast, P22 phage cannot infect this rough strain and its Mu-1 lysogenic derivative WR4027(Mu-1). When a mixture of Salmonella rough specific phage (designated as R phage) was plated on WR4027(Mu-1), smooth derivatives which are resistant to R phage were isolated. Thus, P22 is now able to grow in this new lysogen, WR4027(Mu-1)/R, and give rise to hybrid phages carrying the protein coat of Mu-1 phage and c region of P22. Such hybrids, designated MuimmP22, were isolated by plating the P22 lysates previously grown on WR4027(Mu-1)/R on a smooth but P22-resistant derivative of Mu-1 lysogen, WR4027(Mu-1)/R/22. These results suggest that the Mu-1 tail fibre component of MuimmP22 phage changed its receptor specificity from rough to smooth hosts, indicating inversion of the G region. Thus, MuimmP22 phage infects WR4028 but not WR4027. When P22 high-titer stocks (more than  $10^{10}$  PFU/ml) previously grown on MuimmP22 lysogens of WR4028 were plated on WR4027/R/22(MuimmP22), a few plaques were found. These plaque formers were found to be a new MuimmP22 hybrid phage class which is disimmune over MuimmP22 lysogens. Like λimmP22dis hybrid phage, MuimmP22dis lysogenic derivatives of WR4028 are immune to P22 infection, suggesting that MuimmP22dis carries both c and Im<sup>2</sup> region of P22.

### 2. Isolation of amber suppressor strains of E. coli - S. typhimurium hybrid WR4028/22 met<sup>+</sup>

Both smooth and rough strains of E. coli - S. typhimurium hybrids WR4028 and WR4027 require methionine. We have isolated a met<sup>+</sup> prototroph of WR4028 by

transduction with P22c2 phage. Because of P22c2 infection, all met<sup>+</sup> transductants are rough strains termed WR4028/22 met<sup>+</sup>. Using this prototroph, we have isolated various amino acid requiring mutants. We then looked for suppressor revertants with the consideration that some of the original auxotrophic mutations would be amber mutations. After large scale screening of these revertants by infecting known  $\phi$ 80 amber mutants suppressible by E. coli suppressor I, we found five suppressor mutants. These amber suppressors should be useful for mapping  $\phi$ 80immP22 hybrid phages.

### 3. Isolation and characterization of transducing $\phi$ 80immP22 hybrids

Hybrid  $\phi$ 80immP22 phages, which retain the protein coat of  $\phi$ 80, have been divided into two types with respect to the extent of homology with P22. One hybrid type has a large P22 early gene segment containing the att-erf-c-h21 region. The second type,  $\phi$ 80immP22dis, has a larger P22 segment which includes both immunity (c and immI) regions of P22, i.e., immI-att-erf-c-h21. Since the dis hybrids carry the P22 att region, the prophage is integrated at the P22 insertion site which is near the pro genes of the host. Some of the hybrid phages recovered from lysogens were found to contain reductions in the size of the P22 DNA segment as detected by loss of dis function. In some cases, the total genome length increased despite a reduction in the size of the P22 segment. This increase could represent replacement of a portion of the P22 DNA segment by host chromosomal genes.

Derivatives which have lost the dis function of  $\phi$ 80immP22dis (i.e.,  $\phi$ 80immP22dis<sup>-</sup>) are due to the replacement of the phage segment containing the att through Im genes of P22 with bacterial segment adjacent to the prophage insertion site. As a consequence, the hybrid phage became a high frequency transducing phage for the proline gene but not tryptophan. Since the size of the bacterial segment

substituting for the att-Im segment of the  $\phi 80immP22dis$  hybrid is about equal to that of the  $\phi 80$  inert segment which is about 10% of the  $\phi 80$  genome, the derivative phage should be able to carry a few bacterial genes. Indeed we found that all  $\phi 80immP22dis^-$  carry proA, and argF and about 20% of  $\phi 80immP22dis^-$  isolates also carry metD as demonstrated by cotransduction. Frequency of these transductions are about 10-20% of infected cells for all markers present, indicating that 100% of lysogenized cells are transduced.

#### 4. Complementation of the P22 early gene functions with $\lambda$ and with $\phi 80$

We have previously isolated P22- $\lambda$  hybrid phage which carries a segment of the  $\lambda$  early region and the entire late region of P22. Thus, the  $\lambda$  early genes should regulate expression of the P22 late genes in P22- $\lambda$  hybrid phage. In order to test this possibility, we analyzed complementation capacity of  $\lambda$  or  $\phi 80$  for P22 amber mutants within their early regions. P22 $am24$  mutants spotted on a soft agar overlay containing smooth lysogens carrying either  $\lambda t1tm$  or  $\phi 80$  wild type show replication as observed by formation of plaques or lysis zones whereas P22 $am12$  mutants cannot replicate in these lysogens. Moreover, P22 amber mutants in genes 23 and 7 (endolysin) grow on these lysogens. The same concentration of this P22 amber mutant did not produce any sign of phage replication on a soft agar overlay with a non-lysogenic strain.

#### 5. A new isolation procedure for $\lambda$ -P22Im $\lambda c$ hybrids

P22- $\lambda$  hybrid phage grows on  $\lambda$  lysogens though they share the c immunity region. This is because P22- $\lambda$  hybrid carries the P22 anti-repressor which inactivates  $\lambda$  repressor as well as P22 repressor. When P22- $\lambda$  hybrid phage stocks previously grown on  $\lambda$  lysogens of WR4028 were plated on a  $\lambda$  lysogen of a rough derivative (WR4027), small plaques were found at a frequency of

about  $10^{-5}$ . These plaque formers should be  $\lambda$ -P22Im $\lambda$ c hybrid type because  $\lambda$ immP22 hybrid type carrying  $\lambda$ c and the P22 antirepressor gene can replicate in a rough derivative of  $\lambda$  lysogen, WR4027( $\lambda$ ).

6. Homology between two generalized Salmonella transducing phages ES18 and P22

Salmonella phages have been divided into two subgroups by Boyd: Group A phages are heat stable (resistant to high temperature) and have generalized transducing ability while group B phages are heat labile and do not carry generalized transducing activity with one exception. One of the group B phages, ES18, is a generalized transducing phage. Since we previously found that ES18 is homologous in the c regions, homology analyses were extended to various other genes of P22. Our recent studies show ES18 has homology with the entire early regions of P22. In addition, ES18 and P22 share the generalized transducing gene as observed by complementation of P22am3 mutants with ES18. Thus, we consider that phage ES18 is a generalized transducing hybrid phage as a consequence of recombination between a group A phage and a group B phage.

### Publications

1. Yamamoto, N., Yamamoto, S., Gemski, P., and Baron, L.S., 1981. An unusual  $\lambda$ -P22 phage hybrid with the  $\lambda$   $c^+$  region and the immI region of P22. Abs. Am. Soc. Microbiol. p. 114.
2. Yamamoto, N., Droffner, M.L., Yamamoto, S., Konzelman, J., Gemski, P., and Baron, L.S., 1981. Characterization of hybrids between coliphage  $\phi$ 80 and Salmonella phage P22. Abs. Am. Soc. Microbiol. p. 114.
3. Yamamoto, N., Gemski, P., and Baron, L.S., 1981. Isolation of a hybrid between Salmonella phage P22 and coli mutator phage Mu-1. Abs. Am. Soc. Microbiol. p. 115.
4. Yamamoto, N., Gemski, P., and Baron, L.S., 1982. Genetic studies of new hybrid phage species between coliphage  $\phi$ 80 and Salmonella phage P22. Submitted to J. Gen. Virol.
5. Yamamoto, N., Droffner, M.L., Yamamoto, S., Gemski, P., and Baron, L.S., 1982. Specialized transducing derivatives of  $\phi$ 80immP22 phage. In preparation.

12 Copies

Director (ATTN: SGRD-UWZ-AG)  
Walter Reed Army Institute of  
Research  
Walter Reed Army Medical Center  
Washington, D.C. 20012

4 Copies

HQDA (SGRD-AJ)  
Fort Detrick  
Frederick, Maryland 21701

12 Copies

Defense Documentation Center  
ATTN: DDC-TCA  
Cameron Station  
Alexandria, Virginia 22314

1 Copy

Commandant  
Academy of Health Sciences, U.S. Army  
ATTN: AHS-COM  
Fort Sam Houston, Texas 78234

1 Copy

Dean  
School of Medicine  
Uniformed Services University  
of Health Sciences  
4301 Jones Bridge Road  
Bethesda, Maryland 20014

5 F